

# **NOAA** FISHERIES

NW Fisheries
Science
Center

## NWFSC Fisheries Observer Program Overview

Jon McVeigh
Program Manager,
Fisheries Observation Science (FOS)
September 19th, 2013

### **Presentation Outline**

- Overview of Programs and fisheries observed
- Vessel Selection and Coverage Rates
- Data Collection and QA/QC
- Data Products



#### **Fishery Dependent Data Fishery Data Flows** Commercial **Landings Data Pacific States** (WA, OR, CA) Age-length **Marine Fisheries** Commission **Compositions** Recreational (PacFIN, RecFIN) **Landings Data** (WA, OR, CA) **Total Fishing Observer Data Mortality** (WCGOP, ASHOP) Estimates / **Discard Rates Fishery Independent Data Assessments Shelf-Slope Bottom Trawl Maturity** Survey Data (FRS) **Estimates Data from Other Current** Groundfish Surveys (NWFSC, SWFSC) **Biomass Estimates & Data from Prior Trawl Compositions** Surveys (NWFSC, AFSC) **Hake Biomass** Hake Acoustic & Mid-**Estimates &** water Trawl Data (FEAT) **Compositions**

### GOALS

- Collect high quality, unbiased fisheries-dependent data
- Provide sound science on west coast fisheries bycatch
- Collect and report protected species interactions with west coast fleets



### **Presentation Outline**

- Overview of Programs and fisheries observed
- Vessel Selection and Coverage Rates
- Data Collection and QA/QC
- Data Products



### **Program Information**

#### West Coast Groundfish Observer Program (WCGOP)

- Catch Share Science and catch accounting
- Non-Catch Share Science
- Implemented 2001 with a focus on discards
- Cooperative agreement w/ Pacific States Marine Fisheries Commission (PSMFC)
- 11 NOAA FTEs (3 vacant)
- 14 PSMFC employees

### At-Sea Hake Observer Program (A-SHOP)

- Motherships Science and catch accounting
- Catcher Processers Science and catch accounting
- Implemented in 1977, focus on catch estimation, 2 observers per vessel began in mid-90s



#### Fisheries Observed

Individual Fishing Quota (IFQ)

Fisheries (aka Catch Shares)

Shoreside Hake

Shoreside non-Hake Groundfish

At-sea Hake (3 sub-sectors)

Non-Nearshore Fixed Gear

Limited entry (LE) sablefish

Open Access (OA) fixed gear

Nearshore Fixed Gear

CA and OR

Pink Shrimp Trawl

CA, OR and WA

California Halibut Trawl

#### Gears

**Bottom Trawl** 

**Pelagic Trawl** 

**Shrimp Trawl** 

**Hook and Line** 

Rod and reel

Stick/pipe

Longline

Snap gear

Pots/Traps

Scottish Seine

Kayak



## **Presentation Outline**

- Overview of Programs and fisheries observed
- Vessel Selection and Coverage Rates
- Data Collection and QA/QC
- Data Products



### WCGOP Vessel Selection


### Vessel selections are made by:


- Collecting permit, license, and vessel lists obtained from the NOAA Northwest Region or relevant state agencies' databases.
- Permits/licences/vessels are aggregated by regional "Port Groups" based on previous years landings
- Design-based framework distributes observer coverage more evenly coast-wide than simple random sampling


#### WCGOP Vessel Selection

## Selection period

- 2-month trip period
- LE Trawl fishery (now IFQ) mandatory 100% observer coverage (since 2011)
- Some fisheries are seasonal (LE Fixed Gear SABL, Pink Shrimp)
- Overlapping and stacking permits complicates selection










#### WCGOP Vessel Selection

### **Exceptions:**

- Only vessels >17 feet
- Trips may be waived :
  - Safety
  - Weather
  - Observer availability
  - Vessel space constraints
- No waivers in the IFQ fishery



## **A-SHOP Coverage**

Two observers are deployed on each vessel for every sea day

~100% of hauls are sampled for species

composition




## **Coverage Rates**

**IFQ Shoreside Fishery Coverage Rates, 2011-2012.** Coverage rates are computed as the proportion of total groundfish landings (excluding Pacific hake) from the sampled catch versus total catch (sampled + unsampled). Landed weights were adjusted to fish ticket landing receipts for each trip.

	Bottom Trawl Gears		Hook-and- Line		Pot		Shoreside Hake	
	2011	2012	2011	2012	2011	2012	2011	2012
% tows sampled	99.4%	99.4%	100%	100%	99.9%	99.95%	100%	100%
% groundfish landings sampled	99.4%	99.4%	100%	100%	100%	100%	100%	100%

#### **Non-Catch Share Fisheries**



Observer and Trip Activity for 2002 - 2012. The trip, observer and sea day totals are derived from the fisheries within the respective program. The number of sea days are calculated from the departure and return dates within a trip.

and return dates within a trip.					
PROGRAM	Avg # of Trips per year	Max # of Trips	Min # of Trips		
WCGOP Catch Shares*	2356	2492	2220		
A-SHOP Catch Shares**	41	55	28		
EFP	78	156	29		
Limited Entry	667	818	235		
Open Access	491	717	204		

PROGRAM	Avg # of Observers per year	Max # of Observers	Min # of Observers
WCGOP Catch Shares*	92	97	86
A-SHOP Catch Shares	27	40	16
EFP	8	19	2
Limited Entry	41	46	34
Open Access	35	43	16

<b>└</b> !	U	10	<b>~</b>
Limited Entry	41	46	34
Open Access	35	43	16
PROGRAM	Avg # of Sea Days per year	Max # of Sea Days	Min # of Sea Days
WCGOP Catch Shares*	<del> 7673</del>	7900	7445
A-SHOP Catch Shares	1209	1884	680
EFP	207	556	47
Limited Entry	1874	2515	586
Open Access	801	1395	265
*WCGOP Catch Shares of	data from 2011-2012		
**Trip data only available	for 2008-2012 I.S. Department of Commerce   N		

## **Presentation Outline**

- Overview of Programs and fisheries observed
- Vessel Selection and Coverage Rates
- Data Collection and QA/QC
- Data Products



### **Observers**

### Intensive 3 week observer training

- Observer candidates must meet minimum eligibility requirements such as education standards and required annual briefings
- Covers random sampling, techniques, intense safety training, fish ID, general vessel awareness, conflict resolution and gear use.
- Must pass with 80% or better score on fish and general exams.
- A-SHOP adds additional training for NPGOP certified observers



### **WCGOP Data Collection**

#### Observer weight and sample methods

- Random unbiased sampling
- Multiple sampling methods and priority lists allow flexibility
- Focus on sampling discards and protected species interactions
- Minimum sample sizes, when practical

#### Example weight and sample methods

- Actual weights
- Spatial subsampling (sample a random subsection)
- Basket volume determination
- Tally with collection of subset for average weight



### **WCGOP Data Collection**

#### Trip Level

- Fishing effort
- Total catch estimates
- Fishing locations
- Vessel and crew
- Gear configurations and excluder use

#### Haul/Set Level

- Retained and discarded catch estimates for all catch
- Observer weight methods
- Catch categories system used to estimate single species and species groups



### **WCGOP Data Collection**

#### **Catch Category Level**

- Species composition sampling
- Weights and counts
- Reason for discarding
- Observer sampling methodology

#### **Species Level**

- Lengths, weights, sex and otoliths
- Pacific halibut viabilities
- Genetic samples for salmon and green sturgeon; whole specimens of eulachon (protected species)
- Marine mammal, seabird, sea turtle interactions.



### **A-SHOP Data Collection**

#### Trip

Dates, times, ports, crew size, time lost at sea

#### Haul

- Fishing effort data
- Location, date, time, depths, gear type and performance
- Total catch estimates
- Species composition samples
- Retained and discard estimates
- Biological data on target and bycatch species
- Chinook salmon genetics and coded wire tag sampling
- Protected species interactions and sightings






## **A-SHOP Biological Sampling Summary**


Species	Length	Sex	Weigh t	Age structure	Genetics sample
Hake	X	Х	X	Х	
Bocaccio	X	X	X	Х	
Canary	X	X	X	X	
Darkblotched	X	X	X	X	
POP	X	X	X	Х	
Rougheye	X	X	X	Х	
Widow	X	X	X	X	
Yelloweye	X	X	X	Х	
Yellowtail	X	X	X	X	
Spiny dogfish	X	X	X	Х	
Chinook salmon	X	X	X	Х	X
Sturgeon	X	X	X	Х	Х



## A-SHOP Biological Sampling Summary









#### Yellowtail 7000 1200 otoliths 6000 1000 5000 of lengths or 800 4000 600 3000 Lengths 400 2000 Number Otoliths 200 1000 0 zan zan zan zan zan zan zan zan zan zan

## **A-SHOP Biological Sampling Summary**


#### **Pacific Ocean Perch**




#### Rougheye





#### **Darkblotched**



#### Canary









## WCGOP IFQ QA/QC

- 3 days after landing
- Automated error checks
- In season advising
- Vessel account system 4x/week

Collection and Database entry

Data Submitted to Debriefer

- Trips monthly to debriefer
- Debriefer review
- Forms reviewed
- Accepted/denied

- Feedback to observer
- Corrections within 2 weeks
- Debriefings depending on experience and data quality.

In-Season Feedback

### Data Finalization and Archive

- Error checks (again)
- Data sheets compared against database
- Finalized by debriefer within 2 weeks
- Final data checks in Seattle
- Final data set for the year available by spring of following year.



### A-SHOP Data QA/QC

- Data submitted daily
- In-season data checks
  - Error validations
  - In-season advisor

Deployment (up to 90 days)

End of Deployment /Pre-Debriefing

- Error Validations in NORPAC (Alaska region) database
- Debriefer Review

- Observer interview
- · Data corrected
- 'Debriefed' status

Debriefing

**Data Finalization** 

- Deployment evaluations sent to observers
- Deployment closed



### **Presentation Outline**

- Overview of Programs and fisheries observed
- Vessel Selection and Coverage Rates
- Data Collection and QA/QC
- Data Products



## **Data Processing**

### WCGOP

- Expand species composition by haul
- Translate WCGOP codes to PacFIN codes
- Query and process all PacFIN fish ticket data
- Merge Observer and Fish Ticket data adjust retained catch weights
- Remove non-applicable data: mud, debris/garbage, kelp, rocks etc.



### **Stock Assessment Data Products**

#### **ASHOP** and WCGOP

- Biological Data
  - Lengths
  - Ages
  - Average Weights
- Discard Ratios WCGOP
- % Retained + Total Discard ASHOP
- Total Mortality Estimates
- Discard estimates for fisheries that we observe



### **Stock Assessment Data Products**

### **Groundfish Mortality Estimates**

- Stratification
  - State, latitude, depth, season, fishery, management area
- Discard estimation
- Differs by sector
- General approach: Ratio estimators

Discard (lbs)	X	Fishing Effort Metric	=	Estimated Discard
Fishing Effort Metric	c			
Observed Discard Rate	io	Fleet Total		Fleet Discard

# Strengths

Only source of information for at-sea discard for west coast groundfish species

High (20-99%) coverage levels in the largest harvesting sectors

High retention of observers in non catch share sector

Many products are available online through NWFSC website.



# Strengths

High data quality standards

Comprehensive training program

Frequent feedback and debriefings with observers

Committed staff with extensive observer experience

Internationally recognized safety training program



# Challenges

DATA MANAGEMENT

Data Requests - Confidentiality

Low coverage in some fisheries

Reduced observer numbers (7 of 26 seasonal observers deployed in WCGOP 2013)

Safety of vessels with 100% coverage and aging vessels



# Challenges

New compliance monitoring roles and reporting of near real-time data

Huge increase in data load and observers in the IFQ fisheries

Staffing: 3 vacant positions in an understaffed program

Large geographic area

No dedicated training facility – Train ~150 observers a year

Reliance on AFSC observer program for A-SHOP program



# **Moving Forward**

#### **DATA MANAGEMENT**

- Need data systems to allow for more access and more functions
- Automated system for integrating fisheries dependent data sources

Development of an offline, onboard data entry system

Form collaborative teams for species bycatch reporting

Explore electronic monitoring



# **Moving Forward**

Develop institutional mechanisms to balance priorities with available resources

Develop more non-confidential "off-the-shelf" data summaries

Promote observing as a professional career

Maximize efficiency and reduce uncertainty by exploring alternate sampling strategies and coverage levels





